

## MODEL COMPARISON USING SIMULATED POSTERIOR DRAWS

### 1. THE PROBLEM

- We have two or more models, indexed by  $i$ , each of which, with its prior, defines a joint pdf  $p_i(y, \theta_i)$  for the data and the parameters.
- The posterior probabilities on the models are proportional to  $s_i = \int p_i(y, \theta_i) d\theta_i$ .
- We have no analytic formula for the integrals.

### 2. AN IDENTITY THAT PROVIDES METHODS

If  $q_1(\theta)$  and  $q_2(\theta)$  are positive, integrable functions on the same domain (i.e. can be thought of as unnormalized probability densities), with  $z_i = \int q_i(\theta) d\theta$ , and if  $\alpha(\theta)$  is any function such that  $0 < \int \alpha(\theta) q_i(\theta) < \infty, i = 1, 2$ , then

$$\frac{\int \frac{q_1(\theta) q_2(\theta) \alpha(\theta)}{z_2} d\theta}{\int \frac{q_2(\theta) q_1(\theta) \alpha(\theta)}{z_1} d\theta} = \frac{E_2[q_1 \alpha]}{E_1[q_2 \alpha]} = \frac{z_1}{z_2}.$$

### 3. SPECIFIC METHODS

**importance sampling:**  $\int z_2 = 1, \alpha = 1/q_2, z_1 = E_2[q_1/q_2]$ . Does not use MCMC draws. Blows up if  $q_1/q_2$  is huge for some  $\theta$ 's.

**modified harmonic mean:**  $\int q_2 = 1, \alpha = 1/q_1, z_1 = 1/E_1[q_2/q_1]$ . Uses only MCMC draws. Blows up if  $q_1/q_2$  is huge for some  $\theta$ 's.

**bridge sampling:** Pick  $\alpha$  so both  $q_1 \alpha$  and  $q_2 \alpha$  are bounded, e.g.  $\alpha = 1/(q_1 + q_2)$ . Uses draws from both  $q_1$  and  $q_2$ .

### 4. OPTIMAL $\alpha$

- With same number of draws from  $q_1$  and  $q_2$ , it's

$$\alpha = \frac{1}{z_1 q_2 + z_2 q_1}.$$

- Since we don't know  $z_1/z_2$ , this is not directly a help. But if our initial guess is off, we can update it and repeat — with new  $z_1/z_2$ , but re-using the old draws of  $q_2$  and  $q_1$ .

### 5. WHY "BRIDGE"?

### 6. PATH SAMPLING

---

*Date:* November 16, 2010.

©2010 by Christopher A. Sims. This document may be reproduced for educational and research purposes, so long as the copies contain this notice and are retained for personal use or distributed free.