

# Wold Decomposition

September 21, 2016

## Preliminaries

If we have a set (possibly countably infinite) of random variables  $\{X_i\}$ , the set of all finite linear combinations of them forms a linear space.

We can define an inner product, and thus a norm on that space as  $\langle X, Y \rangle = \text{Cov}(X, Y)$ . Then defining the distance between  $X$  and  $Y$  as  $\|X - Y\|$ , our space is a metric space. We can **complete** the metric space by extending it to include all limits of Cauchy sequences. That is, if  $\{Z_i, i = 1, \dots, \infty\}$  has the property that  $\|Z_m - Z_n\| \rightarrow 0$  as  $m, n \rightarrow \infty$ , then  $Z_\infty = \lim_{i \rightarrow \infty} Z_i$  is in the space.

# Projections

Suppose  $G$  is a complete linear subspace of  $H$ , with a Hilbert space (i.e., innerproduct and norm defined) structure. We can define the operator  $\mathcal{E}$  by

$$\mathcal{E}[X | H] = Z \in H \text{ that minimizes } \|X - Z\| .$$

It is not hard to prove that such a  $Z$  must always exist and is unique.

If  $G_1$  and  $G_2$  are two subspaces of  $H$  such that  $\langle X, Y \rangle = 0$  whenever  $X \in G_1$  and  $Y \in G_2$ , we say that  $G_1$  and  $G_2$  are **orthogonal**, or  $G_1 \perp G_2$  . In that case it is not hard to show that  $\mathcal{E}[X | G_1, G_2] = \mathcal{E}[X | G_1] + \mathcal{E}[X | G_2]$ .

It is always true that  $X - \mathcal{E}[X | G] \perp G$ .

# A finite variance stochastic process and its predictive projections

Now let  $Y_t, t = -\infty, \dots, \infty$  be a vector valued stochastic process. That is, each  $Y_t$  is an  $n$ -dimensional random vector, and the probability law of the stochastic process specifies mutually consistent joint distributions for any finite collection of the  $\{Y_t\}$  variables.

Let  $H_t$  be the complete metric space generated by  $\{Y_s, s \leq t\}$ .

We can always project  $Y_t$  on  $H_{t-1}$  and express the gap between the two as  $\varepsilon_t = Y_t - \mathcal{E}[Y_t | H_{t-1}]$ .

$\varepsilon_t$  is the **innovation** in  $Y_t$ .

## Recursive projections, Wold representation

$H_t$  for any  $t$  is the same as the space spanned by  $\varepsilon_t, H_{t-1}$ . (This is obvious if you think about the definitions.) Therefore we can write

$$y_t = \varepsilon_t + \mathcal{E}[y_t | \varepsilon_{t-1}] + \mathcal{E}[y_t | H_{t-2}] = \varepsilon_t + A_1\varepsilon_{t-1} + \mathcal{E}[y_t | H_{t-2}].$$

The  $A_1$  is a square matrix of coefficients. Since  $\varepsilon_t$  is of dimension  $n$  the space it spans is just the space of linear combinations of elements of the  $\varepsilon_t$  vector, so each element of the  $\mathcal{E}[Y_t | \varepsilon_{t-1}]$  vector is a linear combination of elements of  $\varepsilon_{t-1}$ , given by a row of  $A_1$ .

Repeating this  $T$  times, we get

$$y_t = \sum_{s=0}^{T-1} A_s \varepsilon_{t-s} + \mathcal{E}[Y_t | H_{t-T}] = \tilde{y}_t^T + \bar{y}_t^T.$$

## Taking limits

$\text{Var}(\tilde{y}_t^T)$  is increasing in  $T$  and is bounded above by  $\text{Var}(y_t)$ . (These are matrices, so we mean by “increasing” that their differences are positive semi-definite, which implies their diagonal elements are non-negative.) It is therefore a Cauchy sequence and has a limit we call simply  $\tilde{y}_t$ . This is the **linearly regular** piece of  $y_t$ .

$\text{Var}(\bar{y}_t^T)$  is decreasing in  $T$  and bounded below by zero, so it too is Cauchy and has a limit, which we call  $\bar{y}_t$ .

Note that  $\bar{y}_t$  is in  $H_{t-T}$  for every  $T$ , so  $\mathcal{E}[\bar{y}_t | H_{t-T}] = \bar{y}_t$ , for every  $T$ . In other words,  $\bar{y}_t$  can be forecast arbitrarily well from data on  $y_s$  before time  $t - T$ , no matter how far back in the past these data are. So we call this part the **linearly deterministic** part.

# Stationarity

If the  $y$  process is stationary, meaning the joint distribution of  $\{X_1, \dots, X_m\}$  is the same as that of  $\{X_{s+1}, \dots, X_{s+m}\}$  for any  $s$ , no matter what  $m$  we start with, then our decomposition above produces the same  $A_s$  sequence, no matter what  $t$  we pick for  $y_t$ . Furthermore  $\tilde{y}_t$ ,  $\bar{y}_t$ , and  $\varepsilon_t$  will then also be stationary.

A stationary process is called **linearly regular** if its linearly deterministic component is zero. It is called **linearly deterministic** if its linearly regular component is zero. (For a non-stationary finite variance process, we can do the same decomposition at any  $t$ , but the component processes could have variance zero for some dates and not for others.)

## Examples of LR and LD processes

LR stationary processes: i.i.d.  $N(0, I)$ ; stationary AR(1) ( $\mathcal{E}[y_t | H_{t-1}] = \rho y_{t-1}$ ,  $\text{Var } \varepsilon_t$  constant,  $|\rho| < 1$ )

Examples of LD stationary processes:  $y_t = \sin(t + \theta)$ ,  $\theta \sim U(0, 2\pi)$ ;  $y_t \sim N(0, 1)$ ,  $y_t \equiv y_{t-1}$ ;  $\text{Cov}(y_t, y_{t-s}) = \sin(\pi s/s)$

Note that for the first LD example, making  $\theta$  fixed instead of random, or giving it some other distribution, makes the process non-stationary. This despite the fact that there is no way to tell from observing the time path of  $y_t$  whether  $\theta$  was fixed or random. The third example is one where perfect prediction is not possible with any finite set of past data, but prediction error goes to zero as the amount of past data used goes to infinity.



## Connection to ergodicity

A stationary **ergodic** process  $X_t$  is one such that for any function  $f$  such that  $E[f(X_t)]$  is well defined,

$$\frac{\sum_{t=1}^T f(X_t)}{T} \xrightarrow[T \rightarrow \infty]{a.s.} E[f(X_t)].$$

One is tempted to think of linear regularity as equivalent to ergodicity, but they are not quite the same.

$X_t \equiv X_{t-1}, P[X_t = 1] = .5, P[X_t = 0] = .5$  not ergodic, LD

$X_t = -X_{t-1}, P[X_t = 1] = .5, P[X_t = -1] = .5$  ergodic, LD

$X_t | \sigma^2 \sim N(0, \sigma^2), \text{ i.i.d.}, P[\sigma^2 = 1] = .5, P[\sigma^2 = 2] = .5$  LR, not ergodic